

Calculation of thermodynamic properties of finite Bose-Einstein systems

Peter Borrmann, Jens Harting, Oliver Mülken, and Eberhard R. Hilf

Department of Physics, Carl von Ossietzky University Oldenburg, D-26111 Oldenburg, Germany

(February 1, 2008)

Abstract

We derive an exact recursion formula for the calculation of thermodynamic functions of finite systems obeying Bose-Einstein statistics. The formula is applicable for canonical systems where the particles can be treated as non-interacting in some approximation, e.g. like Bose-Einstein condensates in magnetic traps. The numerical effort of our computation scheme grows only linear with the number of particles.

As an example we calculate the relative ground state fluctuations and specific heats for ideal Bose gases with a finite numbers of particles enclosed in containers of different shapes.

03.75.Fi, 05.30.Jp, 32.80.Pj

With the observation of Bose-Einstein condensation (BEC) of magnetically [1–3] and optically [4] trapped atoms new insights into the nature of this state of matter have been given. The experimental situation is in all cases quite different from the ideal gas treated within the grand canonical ensemble, which is the standard textbook example. First, the number of particles within the traps is fixed and finite, which suggests a canonical or micro-canonical treatment of the systems. Second, the confining trap potentials greatly influence the condensate properties. Third, although the trapped gases are quite dilute the validity of the treatment as non-interacting particle gases has to be checked from case to case.

Even within the approximation of non-interacting particles the calculation of the thermodynamic properties of the Bose-Einstein systems remains a hard to tackle mathematical problem. Recently, some approximate methods to calculate the fluctuation of the ground state occupation number in a trapped Bose-Einstein condensate have been developed [5–8]. Here we present an exact method to calculate all thermodynamic quantities of finite canonical Bose systems, given the one particle density of states.

As the starting point we utilize the recursive formula of the canonical partition function for a system of N noninteracting bosons as given in [9]:

$$Z_N(\beta) = \frac{1}{N} \sum_{k=1}^N Q_k(\beta) Z_{N-k}(\beta) , \quad (1)$$

where $Q_k(\beta) = Z_1(k\beta) = \sum_i \exp(-k\beta\epsilon_i)$ is the one-particle partition function at the temperatures $k\beta$ and $Z_0(\beta) = 1$. The microcanonical partition $\Gamma_N(E)$ can be calculated by an inverse Laplace-transform of (1) and is given by

$$\begin{aligned} \Gamma_N(E) &= \frac{1}{N} \sum_{k=1}^N \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} d\beta \exp(\beta E) Q_k(\beta) Z_{N-k}(\beta) \\ &= \frac{1}{N} \sum_{k=1}^N \int_0^E dE' \Gamma_1^k(E') \Gamma_{N-k}(E - E') , \end{aligned} \quad (2)$$

where $\Gamma_1^k(E)$ is the inverse Laplace-transform of $Q_k(\beta)$ and $\Gamma_0(E) = \delta(E)$. A similar, slightly less general, equation has recently been derived by Weiss and Wilkens [10].

Eq.(1) can be used to calculate all thermodynamic quantities by appropriate differentiation of $\ln Z_N$. However, in any case Z_N occurs as a normalization factor and has to be

calculated explicitly. This turns out to be a major drawback. First, the numerical effort to calculate Z_N grows with the square of the particle number N . Moreover, since $Z_N(\beta)$ grows exponentially with N multiple precision arithmetic is required for proper calculation. We will present a method avoiding these difficulties.

To ease our derivations we rewrite $Z_N(\beta)$ utilizing the $\mathcal{Z}$ -transform and define:

$$\mathcal{Z}(Z) = F(x) = \sum_{k=0}^{\infty} \frac{Z_k(\beta)}{x^k}, \quad (3)$$

$$\mathcal{Z}(Q) = G(x) = \sum_{k=0}^{\infty} \frac{Q_k(\beta)}{x^k}, \quad (4)$$

where we define $Q_0(\beta) = 0$. Taking advantage of the basic properties of the $\mathcal{Z}$ -transform eq.(1) can be written in the form

$$-x \frac{d}{dx} F(x) = F(x)G(x), \quad (5)$$

yielding ¹

$$F(x) = \exp \left(\sum_{k=1}^{\infty} \frac{Q_k(\beta)}{k} x^{-k} \right). \quad (6)$$

Applying the inverse $\mathcal{Z}$ -transform we may write $Z_N(\beta)$ as

$$Z_N(\beta) = \frac{1}{2\pi i} \int_C F(x) x^{N-1} dx, \quad (7)$$

where $C := \{z \in \mathbb{C} : |z| = r\}$ and r has to satisfy the condition $|Z_N(\beta)| \leq \exp(rN)$.

Alternatively we may write:

$$Z_N(\beta) = \frac{1}{N!} \left. \frac{d^N}{dx^N} F(1/x) \right|_{x=0}. \quad (8)$$

Using eq.(8) the number of particles with energy ϵ_i can be calculated by

$$\begin{aligned} \eta_i(N, \beta) &= \frac{-1}{\beta} \frac{\partial}{\partial \epsilon_i} \ln Z_N(\beta) \\ &= \frac{1}{Z_N(\beta)} \sum_{k=1}^N \exp(-\beta k \epsilon_i) Z_{N-k}(\beta) \end{aligned} \quad (9)$$

¹Note that $F(x)$ is closely related to the grand canonical partition function.

Some reordering yields:

$$\eta_i(N+1, \beta) = \frac{Z_N(\beta)}{Z_{N+1}(\beta)} \exp(-\beta\epsilon_i)(\eta_i(N, \beta) + 1) . \quad (10)$$

Since the particle number is a conserved quantity in the canonical ensemble the direct calculation of the normalization factor can be omitted by using the relation

$$\frac{Z_N(\beta)}{Z_{N+1}(\beta)} = \frac{N+1}{\sum_{i=0}^{\infty} \exp(-\beta\epsilon_i)(\eta_i(N, \beta) + 1)} . \quad (11)$$

For Fermi systems the following recursion formula

$$\eta_i(N+1, \beta) = \frac{Z_N(\beta)}{Z_{N+1}(\beta)} \exp(-\beta\epsilon_i)(1 - \eta_i(N, \beta)) , \quad (12)$$

with

$$\frac{Z_N(\beta)}{Z_{N+1}(\beta)} = \frac{N+1}{\sum_{i=0}^{\infty} \exp(-\beta\epsilon_i)(1 - \eta_i(N, \beta))} . \quad (13)$$

can be derived in a similar manner. In practice, only a limited number of energy levels has to be taken into account, since the occupation probability rapidly decreases with increasing energy eigenvalues. Equations (10) and (11) are extremely useful in practical calculations. The numerical effort to calculate the occupation numbers grows only linear with the number of particles. Moreover, only a moderate arithmetic precision is required. Having the occupation probabilities at hand the energy expectation value is given by:

$$E(N, \beta) = \sum_{i=0}^{\infty} \epsilon_i \eta_i(N, \beta) . \quad (14)$$

The calculation of the fluctuation of the occupation probabilities $\delta\eta_i(N, \beta)$ is a little bit more complicated and contains another recursion:

$$\begin{aligned} (\delta\eta_i(N+1, \beta))^2 &= \frac{1}{\beta^2} \frac{\partial^2}{\partial \epsilon_i^2} \ln(Z_{N+1}(\beta)) \\ &= \frac{-1}{\beta} \frac{\partial}{\partial \epsilon_i} \eta_i(N+1, \beta) \\ &= \frac{Z_N(\beta)}{Z_{N+1}(\beta)} \exp(-\beta\epsilon_i) \left\{ \delta^2 \eta_i(N, \beta) \right. \\ &\quad \left. + [\eta_i(N+1, \beta) + 1] [\eta_i(N, \beta) - \eta_i(N+1, \beta) + 1] \right\} . \end{aligned} \quad (15)$$

To illustrate the usefulness of our recursion formulas we consider the ideal gas with parameters of liquid Helium in containers of different shapes :

i) a cube with side-length L_x, L_y, L_z , and energy levels

$$E_{n_x, n_y, n_z} = \frac{\pi^2 \hbar^2}{2m_{He}} \left(\frac{n_x^2}{L_x^2} + \frac{n_y^2}{L_y^2} + \frac{n_z^2}{L_z^2} \right) \quad (16)$$

ii) a sphere with radius a and energy levels

$$E_{n,l} = \frac{\hbar^2}{2m_{He}a^2} u_{n,l} \quad (17)$$

and degeneracy $\sigma_{n,l} = 2l + 1$, and

iii) a cylinder with diameter $d = 2a$, height L , energy levels

$$E_{n,l,m} = \frac{\hbar^2}{2m_{He}} \left(\frac{v_{n,l}^2}{a^2} + \frac{m^2 \pi^2}{L^2} \right) \quad (18)$$

with $n = 1, 2, \dots$, $l = 1, 2, 3, \dots$, and $m = \dots, -1, 0, 1, \dots$

We denoted the zeros of the half integer Bessel functions $J_{n+1/2}(r)$ by $u_{n,l}$ and the zeros of the integer Bessel function $J_n(r)$ by $v_{n,l}$.

Fig. 1 displays the specific heats and the fluctuations of the ground state occupation number $\delta\eta_0/N$ as a function of the canonical temperature for different trap geometries and $N = 100, 1000$, and 10000 He-atoms. In all cases the particle density is taken to be $\rho = 0.0216 \text{\AA}^{-3}$. With growing system size the differences between the specific heats for the different trap geometries almost vanish and approach the typical shape of the curve for the ideal Bose gas. I.e., with respect to the specific heat the boundary conditions get more and more unimportant with increasing volume. In contrast, the ground state fluctuations exhibit a completely different behaviour. The cubic box, the *compact* cylinder with equal diameter and height, and the sphere show almost equal ground state fluctuations for all system sizes, while the ground state fluctuations of the stretched box and the stretched cylinder are remarkably larger for temperatures below the critical temperature. This effect is not unexpected because restricting the particle motion in one or two dimensions makes the system act like a lower dimensional system, which are known to show larger fluctuations. Moreover, the differences between the fluctuations of the *stretched* traps and the *compact* traps

do not decrease with increasing system size. The reason for this behaviour is found in the energy difference between the ground state and the first excited level, which is much larger for the *stretched* traps than for the *compact* traps. Since $\delta\eta_0/N$ decreases approximately with $N^{-1/3}$ the infinite particle number limit is the same for all trap geometries. Under experimental considerations our results imply that the stability of the condensate fraction in anisotropic traps should be considerably smaller than in isotropic traps. In Fig. 1 we plotted $\delta\eta_0/N$ to allow good comparison with previous published results [8,5]. Since this quantity goes to zero as the system size increases it is a bad indicator for phase transitions. The relative ground state fluctuation $\delta\eta_0/\eta_0$ shown in Fig. 2 is much more conclusive in this respect.

True phase transition only occur for infinite systems. However, it is well known from other systems, e.g. finite spin lattices and clusters [11,12], that finite systems already display the onset of phase transitions. Instead of having a well defined critical temperature the transition occurs in a broader crossover region. As can be extracted from Fig. 2 the crossover region, which is indicated by the sharp increase of $\delta\eta_0/\eta_0$, extends even for the cubic box with 10000 atoms over a temperature range of 0.5 K. For the *stretched* trap the crossover region is about twice as large.

All calculations have been performed on an IBM-43P (233 MHz) workstation. For 1000 particles a run with 200 temperature points took about six minutes, for 10000 particles one hour. A calculation of the same quantities with the recursion formula given in equation (1) takes for 1000 particles about six hours (and would take at least 600 hours for 10000 particles). Moreover, due to the numerical instabilities connected with (1) $N = 1000$ was the largest particle number we achieved with this formula, even though we utilized a multiple precision package.

We expect the recursion formulas to be quite useful for calculating the properties of dilute atomic Bose gases in magnetic traps with different geometry. For this propose we provide

an easy to use Java program², which requires as input only the energy level distribution with appropriate degeneracy and calculates the basic properties of finite Bose systems with particle numbers up to 10000. A slightly faster Fortran code is available upon Email request³.

²The code is available within the World-Wide-Web:

<http://www.physik.uni-oldenburg.de/~borrmann/BEC>

³Corresponding author: *borrmann@uni-oldenburg.de*

REFERENCES

- [1] C. Bradley, C. Sackett, J. Tollet, and R. Hulet, Phys. Rev. Lett. **75**, 1687 (1995).
- [2] M. Anderson *et al.*, Science **69**, 198 (1995).
- [3] K. Davies *et al.*, Phys. Rev. Lett. **75**, 3969 (1995).
- [4] D. Stamer-Kurn *et al.*, Phys. Rev. Lett. **80**, 2027 (1998).
- [5] S. Grossmann and M. Holthaus, Phys. Rev. Lett. **79**, 3557 (1997).
- [6] S. Grossmann and M. Holthaus, OPTICS EXPRESS **1**, 262 (1997).
- [7] P. Navez *et al.*, Phys. Rev. Lett. **79**, 1789 (1997).
- [8] M. Gajda and K. Rzazewski, Phys. Rev. Lett. **78**, 2686 (1997).
- [9] P. Borrmann and G. Franke, J. Chem. Phys **98**, 2484 (1993).
- [10] C. Weiss and M. Wilkens, OPTICS EXPRESS **1**, 272 (1997).
- [11] R. Kunz and R. Berry, Phys.Rev.Lett. **71**, 3987 (1993).
- [12] D. Wales and R. Berry, Phys.Rev.Lett. **73**, 2875 (1994).

FIGURES

FIG. 1. (a-c) Specific heats C_v/N and (d-f) ground state fluctuations as a function of the canonical temperature for systems of $N = 100$, $N = 1000$, and $N = 10000$ particles. The solid lines represent the results for a spherical trap, the dashed lines for a cube, the circles for a cylinder with a diameter to height ratio of $d/L = 1$, the long dashed lines for a box with side-lengths $L_z = 4L_y = 4L_x$, and the squares for a cylinder with $d/L = 1/4$. In all cases the particle density is taken to be $\rho = 0.0216 \text{\AA}^{-3}$.

FIG. 2. Relative ground state fluctuations $\delta\eta_0/\eta_0$ as a function of temperature for the cubic box and the stretched box and $N = 1000$, and 10000 particles.



